

Clinician Update

Pericarditis

Cardiology



Criteria for diagnosis

P. 2

Testing

P. 2

End notes

P. 3

DIAGNOSIS

CRITERIA

Because it is not always obvious what is causing someone's chest pain after they present to the ER, doctors use criteria to try to be more certain that a patient has pericarditis.

LAB

Signs of inflammation — Since pericarditis is an inflammatory disease, laboratory signs of inflammation are common in patients with acute pericarditis. These include elevations in the white blood cell count, erythrocyte sedimentation rate, and serum C-reactive protein concentration. While elevation in these markers supports the diagnosis, they are neither sensitive nor specific for acute pericarditis. Additionally, in the hyperacute phase of pericarditis, these markers may remain normal, and increased levels may be found only on follow-up. However, markers of inflammation play a role in determining the optimal duration and approach to tapering of therapy.

Acute pericarditis (at least 2 criteria of 4 should be present):


Typical chest pain

Pericardial friction rub

Suggestive ECG changes (typically widespread ST-segment elevation)

New or worsening pericardial effusion

ECG



TREATMENT

MEDICAL

- *NSAID*
- *glucocorticoid*
- *Colchicine*
- *IL1 inhibitors*

NSAID

Glucocorticoid

Start at 0.5 mg per kg orally and reduce by 10 mg. Every week as symptoms tolerate

Colchicine

IL1 inhibitor

Anakinra (Kineret)

Dose is 100 mg. SC daily. Cost is expensive. Requires specialty pharmacy and prior authorization.



TERMINOLOGY

INCESSANT

Incessant pericarditis is recurring or unable to discontinue acute treatments without a relapse.

REFERENCES

1. M Imazio. Acute pericarditis: Clinical presentation and diagnosis. Uptodate.